

Productive indicators in *Cavia porcellus*, fed five forage species in the Austro region of Ecuador[▲]

Wilson Geovanny Macancela-Urdiales¹, <https://orcid.org/0000-0002-5501-1087>, Mildrey Soca-Pérez², <https://orcid.org/0000-0002-8962-9993> and Tania Sánchez-Santana², <https://orcid.org/0000-0002-2634-830X>

¹Propietario de la finca "Gupancay". Parroquia de San Juan del Cid, cantón Gualaceo. Provincia del Azuay, Ecuador. ²Estación Experimental de Pastos y Forrajes Indio Hatuey, Universidad de Matanzas, Ministerio de Educación Superior. Central España Republicana, C.P. 44280, Matanzas, Cuba. E-mail: whatmons@yahoo.es

Abstract

Objective: To evaluate the performance of productive indicators in *Cavia porcellus* (Guinea pig), fed five forage species, during the growth-fattening stage.

Materials and Methods: A study was conducted in the Gupancay farm, located in the Gualaceo canton, Azuay province, Ecuador. A complete randomized experimental design was applied with five treatments, which corresponded to the evaluated species: T1- *Medicago sativa* L., T2- *Alnus acuminata* Kunth, T3- *Cenchrus purpureus* (Schumach.) cv. Morado, T4- *Tithonia diversifolia* (Hemsl.) A. Gray and T5= *Saccharum officinarum* L. + balanced feed. The duration of the experimental period was four months. The diets were formulated with the forage species and the concentrate feed, in 70-30 % ratio for all the treatments. Fifteen male crossbred newly-weaned Guinea pigs, similar in age and weight, were used. The evaluated indicators were: live weight, weight gain, feed conversion and carcass yield.

Results: The best bromatological composition was for *M. sativa* and *T. diversifolia*, with crude protein values of 20 and 21 %, respectively. *A. acuminata* showed the lowest protein values (5 %), but the highest crude fiber (35,2 %) and organic matter ones (94,8 %). However, the *in vitro* organic matter digestibility was low with regards to the other forage plants (59,3 %). The best productive results were for the treatments that used *M. sativa* and *T. diversifolia*. The lowest values were recorded in *S. officinarum* and *A. acuminata*, with significant differences ($p < 0,001$).

Conclusions: The best performance for the variables weight gain, feed conversion and carcass yield were obtained in the treatments that included *M. sativa* and *T. diversifolia* in the diet of Guinea pigs. These forage species had a crude protein percentage higher than 20 %.

Keywords: *Tithonia diversifolia*, Guinea pigs, *Medicago sativa*

Introduction

The production of Guinea pigs in Ecuador is, generally, a rural activity, in which the traditional-family rearing system prevails. The Guinea pig constitutes one of the most traditional dishes of the Ecuadorian Andes and its elaboration represents an employment source for hundreds of farmers. Hence it is considered by the UN and FAO as a source of food security for the world population with scarce economic resources (Rosales Jaramill *et al.*, 2018).

Guinea pigs are monogastric, herbivore animals. Their digestive physiology is characterized by post-gastric cecal fermentation (Canto and Bernal-Wilmer, 2018). Diverse research works have been conducted about Guinea pig feeding (Choez and Ravillet, 2018), in order to evaluate its effect on the productive parameters (pregnancy and parturition rate, litter size, sow and pup morbidity, post-parturition weight of the sows, growth rate for

fattening, weight at slaughter and carcass quality, among others).

Fresh alfalfa (*Medicago sativa* L.), barley (*Hordeum vulgare* L.), oat (*Avena sativa* L.) and wheat (*Triticum vulgare* L.) constitute the principal basis of this mammal feeding, because they contribute important energy and protein quantities to the diet (Alvarado-Zuta, 2017). Different research works have aimed at obtaining adequate feedstuffs for the nutritional requirements of these animals in order to achieve optimum production levels. The use of grasses and tropical shrub forage plants (Meza-Bone *et al.*, 2014), dry velvet bean forage [*Stizolobium deeringianum* L. Medik.] (Valenzuela-Rocha, 2015), roots and foliage of cassava [*Manihot esculenta* Crantz] (Paillacho-Sánchez, 2017), as well as the inclusion of cowpea (*Vigna unguiculata* L. Walp) as ingredient in rations for growth-fattening (Choez and Ravillet, 2018), are among the feeding

[▲]Paper presented in the 5th International Convention Agrodesarrollo 2019 celebrated on October 22-26, 2019. Plaza America Convention Center. Varadero, Cuba

Received: July 11, 2019

Accepted: September 20, 2019

How to quote a paper: Macancela-Urdiales, W. G.; Soca-Pérez, Mildrey & Sánchez-Santana, Tania. Productive indicators in *Cavia porcellus*, fed five forage species in the Austro region of Ecuador. *Pastos y Forrajes*. 42 (4):246-251, 2019.

This is an open access article distributed in Attribution NonCommercial 4.0 International (CC BY-NC4.0) <https://creativecommons.org/licenses/by-nc/4.0/>
The use, distribution or reproduction is allowed citing the original source and authors.

sources that can cover the nutritional needs of this species. The use of different proportions of cattle blood meal (Bazán-García *et al.*, 2016) and the stubble of such plants or crops as pepperweed [*Lepidium peruvianum* Chacón] (Castro-Bedriñana *et al.*, 2018), among other feedstuffs, is also included.

The feeding and nutrition needs of Guinea pigs vary according to the stages of their life cycle (lactation, growth and reproduction). However, in all of them protein, energy, fiber, vitamins, minerals and water are required. Hence the production and profitability of Guinea pig rearing will depend on the feeding sources (Castro-Bedriñana *et al.*, 2018).

From the above-explained facts, the objective of this research was to evaluate the performance of productive indicators in *Cavia porcellus* (Guinea pig), fed five forage species, during the growth-fattening stage.

Materials and Methods

Characteristics of the research area. The study was conducted in the Gupancay farm, owned by the Macancela-Urdiales family, located on the old road to San Juan del Cid, km 6^{1/2}, in the San Juan parish, Gualaceo canton, Azuay province, Ecuador. The farm has a total area of 9 ha and is located in a zone of mountainous topography, at an altitude of 2 400 m.a.s.l., on a Vertisol soil. The mean annual rainfall is 1 100 mm, distributed throughout the year. The mean temperature is 17 °C (table 1).

Experimental design and treatments. A complete randomized design was used with five treatments and 15 animals per treatment, each one was considered a repetition. The duration of the experimental period was four months (16 weeks) and was in correspondence with the growth-fattening stage.

T1- *Medicago sativa* L. + balanced feed

T2- *Alnus acuminata* Kunth + balanced feed

T3- *Cenchrus purpureus* (Schumach.) cv. Morado + balanced feed

T4- *Tithonia diversifolia* (Hemsl.) A. Gray + balanced feed

T5- *Saccharum officinarum* L. + balanced feed

Characteristics and management of the forage species. Each species was managed according to the crop characteristics:

- *M. sativa* (alfalfa). Legume, cutting frequency of the forage between 50 and 60 days.
- *A. acuminata*. Betulaceae, tree, three years as average after being planted.
- *C. purpureus* cv. Morado (king grass morado). Grass, cutting frequency of the forage between 50 and 60 days.
- *T. diversifolia* (Mexican sunflower). Asteraceae, cutting frequency of the forage between 50 and 60 days.
- *S. officinarum* (sugarcane). Gramineae, plantation with two years after the establishment, without homogenization cut.

The forages were cut in the morning and were chopped with a chopping machine F-150 (4 mm cutting size) to homogenize the feedstuffs.

Animal management and feeding. Male, crossbred, weaned Guinea pigs were used, similar in age (15 ± 3 days) and weight (381 g). They were identified with aluminum ear rings and randomly distributed to form homogeneous groups. They received a diet constituted by forages (according to the treatment) and concentrate feed. The balanced feed was offered in the morning; while the forage species were chopped and distributed at three moments of the day (morning, afternoon, night). The animals were subject to an adaptation period of 15 days, before the experimental period.

Measurements. Of each forage species, three 500-g samples were taken during the experimental period. They were transferred to the laboratory of chemical sciences of the University of Cuenca in order to determine their bromatological composition, according to the technique proposed by the AOAC (2016): dry matter (DM), organic matter (OM), crude protein (CP), crude fiber (CF), ash (ASH), calcium (Ca), magnesium (Mg) and phosphorus (P). For determining the organic matter *in vitro* digestibility (OMD), the KOH technique was used.

Table 1. Climate data of the experimental stage.

| Variable | September | October | November | December | January | February |
|--|-----------|---------|----------|----------|---------|----------|
| Mean temperature, °C | 16,6 | 17,3 | 17,6 | 17,6 | 17,5 | 17,6 |
| Mean temperature of the last ten years, °C | 16,6 | 17,3 | 17,6 | 17,6 | 17,5 | 17,6 |
| Mean rainfall, mm | 60 | 75 | 75 | 70 | 65 | 80 |
| Mean rainfall of the last ten years, mm | 60 | 77 | 77 | 74 | 63 | 81 |

Source: Provincial Government of Azuay (2015)

Feed intake. It was determined through an offer-consumption-refuse test (determined by the difference of the supplied feed during the day and the waste recorded the next day) with a weekly frequency.

Live weight and weight gain. The animals were weighed with a monthly frequency. A Century® digital scale was used, with maximum capacity of 10 kg and accuracy of $\pm 0,5$ kg. The accumulated weight gain for the period was quantified. The mean daily gain (MDG) was expressed in grams/animal/day, from the differences of initial and final live weight.

Feed conversion (FC). It was calculated establishing the relation between the feed intake and weight gain, according to the formula:

$$FC = \frac{\text{consumed feed (kg/animal/period)}}{\text{live weight gain, kg}}$$

Carcass yield (CY). It was determined by considering the weight of the warm carcass weight with regards to the live weight of the animal at the moment of slaughter. Head, legs and viscera (heart, liver and kidney) were included. The following formula was used:

$$CY = \frac{\text{carcass weight} \times 100}{\text{live weight gain, kg}}$$

Statistical analysis. The results were recorded in an Excel database. Simple variance analysis was used, after testing the variance homogeneity (Levene's test) and normal distribution (Shapiro-Wilk test) assumptions. For the mean comparison, Tukey's test was used for a significance level of $p < 0,05$ through the statistical package SSPS®, version 10.0.1 for Windows.

Results and Discussion

Table 2 shows the bromatological composition of the evaluated forage species. The DM, CP and CF values are in correspondence with the bromatological indicators described for *M. Sativa* (Aguilar-Quintana 2017), *Saccharum spp.* (Salazar-Ortiz *et al.*, 2017), *T. diversifolia* (Rivera *et al.*, 2019) and *C. purpureum* (Vivas-Quila *et al.*, 2019).

The best bromatological composition was for *M. sativa* and *T. diversifolia*, with CP values of 20 and 21 %, respectively. Both species represent a potential for Guinea pig feeding, when considering that this indicator is one of the most important, because their nutritional requirements can vary between 14 and 22 % depending on the phase of their productive cycle (pregnancy, lactation, growth), as stated by señala (Alvarado-Zuta, 2017)

A. acuminata showed the lowest protein values and the highest CF ad OM; while the *in vitro* OM digestibility was low with regards to the other forage plants (59,3 %).

The highest ASH, Ca, P and Mg values were for *T. diversifolia* and *M. sativa*. According to Gualoto-Lata (2018), minerals fulfill multiple functions in the organism, among them the structural, physiological and catalyst ones, the most important ones being the formation of bones and nerves.

Table 3 shows the initial and final live weight, total weight gain and feed conversion. The best productive results were for the treatments that used *M. sativa* and *T. diversifolia*. The lowest values were recorded in *S. officinarum* and *A. acuminata*, with significant differences ($p < 0,001$).

The values of final live weight in the forage species oscillated between 1 100 and 1 500 g/animal. These are similar to the ones reported by Camino and Hidalgo (2014), who evaluated a mixed diet based on balanced feed, vitamin C (70 mg/100 g of feedstuff) and broccoli (*Brassica oleracea* L.) stubble. Nevertheless, they were higher than the ones reported by Andrade-Yucailla *et al.*, (2015), when evaluating three substitution levels (25, 40 and 55 %) of green forage of *Ipomoea batatas* L. (sweet potato) in diets for Guinea pigs during the growth-fattening stages in the Ecuadorian Amazonia region. These authors reported values between 910 and 1 050 g/Guinea pig.

The results of final live weight for the treatment with *S. officinarum* were among the lowest ones, but were higher than those referred by Toro-Molina *et al.* (2017), who used 15 % of bagasse of this plant

Table 2. Bromatological composition of the forage species, %.

| Treatment | DM | CP | CF | OM | ASH | Ca | Mg | P | OMD |
|------------------------|----|----|------|------|-------|-----|-----|-----|------|
| <i>M. sativa</i> | 23 | 20 | 24,4 | 88,0 | 12,00 | 5,2 | 2,7 | 0,6 | 72,4 |
| <i>A. acuminata</i> | 37 | 5 | 35,2 | 94,8 | 5,16 | 2,3 | 1,9 | 0,4 | 59,3 |
| <i>C. purpureus</i> | 16 | 10 | 28,8 | 82,6 | 17,40 | 2,2 | 1,7 | 0,4 | 65,2 |
| <i>T. diversifolia</i> | 20 | 21 | 25,1 | 84,4 | 15,59 | 5,3 | 2,6 | 0,7 | 68,6 |
| <i>S. officinarum</i> | 36 | 6 | 24,8 | 97,2 | 2,76 | 2,4 | 1,7 | 0,3 | 67,8 |

Table 3. Performance of the productive indicators of Guinea pigs fed five forage species.

| Treatment | Initial weight, g | Final weight, g | Total weight gain, g | Feed conversion |
|------------------------|-------------------|------------------------------|-----------------------------|--------------------------|
| <i>M. sativa</i> | 386,9 ± 21,65 | 1427,0 ± 45,92 ^a | 1040,1 ± 59,83 ^a | 8,8 ± 0,49 ^a |
| <i>A. acuminata</i> | 393,4 ± 26,24 | 1130,5 ± 37,39 ^b | 737,1 ± 40,22 ^b | 15,8 ± 0,95 ^b |
| <i>C. purpureus</i> | 367,0 ± 19,62 | 1239,8 ± 48,54 ^{ab} | 872,8 ± 47,46 ^{ab} | 9,8 ± 0,51 ^a |
| <i>T. diversifolia</i> | 375,3 ± 18,23 | 1262,5 ± 33,73 ^{ab} | 887,3 ± 41,84 ^a | 8,5 ± 0,40 ^a |
| <i>S. officinarum</i> | 381,5 ± 15,16 | 1146,5 ± 34,38 ^b | 765,0 ± 39,78 ^b | 13,9 ± 0,69 ^b |
| P-value | 0,891 | 0,000 | 0,000 | 0,000 |

a, b: Different letters in the same column indicate significant differences ($p \leq 0,05$)

(935,2 g). They were also better than those obtained by Avalos-Sánchez (2010), when evaluating four inclusion levels of chopped fresh sugarcane (20, 40, 60 and 80 %), in both cases in mixed diets with *M. sativa* during growth-fattening.

The values recorded in this research for total weight gain in the period were higher than the ones reported by Sánchez-Laiño *et al.* (2009), who evaluated three tropical grasses (*Panicum maximum*¹ Jack, *Zea mays*, *S. officinarum*) plus concentrate feed during the fattening of improved Guinea pigs in the Ecuadorian coastal region. These authors obtained values of 478,5; 521,9 and 398,3 g/period, respectively.

The MDG of the animals was similar to the one reported by Meza-Bone *et al.*, (2014), when evaluating forage tropical grasses (*P. maximum*; *Pennisetum* sp.) and shrubs (*Morus alba* L., *Erythrina poeppigiana* (Walp.) O.F.Cook, *Hibiscus rosa-sinensis* L.) *ad libitum* in Guinea pig feeding in Quevedo, Ecuador. However, they were higher than the 10,5 g/animal/day referred by Yamada *et al.* (2019) in the evaluation of two meat lines of Guinea pigs (improved genotypes) in the central coast of Peru, with feeding regime based on forage corn, wheat bran and fresh water.

Significant differences were found ($p > 0,001$) in the average feed conversion among the treatments *M. sativa*, *T. diversifolia* and *C. purpureus* with regards to *S. officinarum* and *A. acuminata* (table 3). These results are similar to the ones reported by Guevara and Carcelén (2014) when evaluating the effect of probiotics on improved Guinea pigs for the growth and fattening stage. They also coincide with those referred in the study conducted by Andrade-Yucailla *et al.* (2015) about the evaluation of grasses adapted to the Amazonian region.

When evaluating the carcass yield (table 4), the values were between 71 and 76 %. The best results

were found in *M. sativa* and *T. diversifolia*. They were followed by *C. purpureus* and *S. officinarum*. In studies conducted by Camino and Hidalgo (2014), these authors obtained carcass yields similar to the ones recorded in this work, with values of 72 and 73 %, when evaluating different feeding alternatives (concentrate feed as pellet and meal and concentrate feeds with restriction of the forage offer, respectively). However, the result of this study slightly surpassed those obtained by Canto *et al.*, (2018), who evaluated the effect of supplementation with probiotic (*Lactobacillus*) in diets of *M. sativa* and concentrate feed, and obtained values between 70 and 72 %.

Table 4. Carcass yield of the animals for the evaluated forage species.

| Treatment | Carcass yield,% |
|------------------------|-----------------|
| <i>M. sativa</i> | 75 |
| <i>A. acuminata</i> | 72 |
| <i>C. purpureus</i> | 74 |
| <i>T. diversifolia</i> | 75 |
| <i>S. officinarum</i> | 74 |

The animals that consumed the *T. diversifolia* forage showed higher efficiency in their conversion from feed to meat, without significant differences regarding the carcass yield, for which it could be inferred that this species has the same nutritional effect as *M. sativa*, because significant differences were not found either between both treatments for the live weight and animal gain.

The productive indicators were very affected in *A. acuminata*, if compared with the other forage plants. This performance could have been influenced by the lower percentages in forage utilization, as well as the low protein contents and high fiber levels, characteristics which qualify *A. acuminata* as a local resource with limitations for Guinea pig feeding.

¹Current scientific name: *Megathyrsus maximus* (Jacq.) B.K.Simon & S.W.L.Jacobs

Conclusions

The best performance for the variables weight gain, feed conversion and carcass yield were obtained in the treatments that included *M. sativa* and *T. diversifolia* in the diet of Guinea pigs. These forage species had a crude protein percentage over 20 %.

Acknowledgements

The authors thank the Macancela Urdiales family for the funding for this research. Thanks are also expressed to the Austro Research Station of the National Institute of Agricultural Research (INIAP for its initials in Spanish), especially to Eng. Maximiliano Ochoa and to the School of Agricultural Sciences of the University of Cuenca for the support to conduct this study.

Authors' contribution

- Wilson Geovanny Macancela-Urdiales. Conducted the experiments and data collection. In addition, he prepared the work for its publication.
- Mildrey Soca-Pérez. Conceptualized the research idea and supervised the research activity.
- Tania Sánchez-Santana. Carried out the statistical analysis and reviewed the manuscript.

Conflicts of interests

The authors declare that there are no conflicts of interests.

Referencias bibliográficas

- Aguilar-Quintana, E. D. *Producción de biomasa forrajera de variedades o ecotipos de alfalfa (Medicago sativa L.) en el sector humedades del distrito de Salas-Lambayeque*. Tesis presentada como requisito para optar el título profesional de Ingeniero Zootecnista. Lambayeque, Perú: Universidad Nacional Pedro Ruiz Gallo, 2017.
- Alvarado-Zuta, R. *Efecto de la yaca (Artocarpus heterophyllus) y alfalfa (Medicago sativa) en el comportamiento productivo de cuyes (Cavia porcellus) en fase de crecimiento y engorde*. Tesis para optar el título profesional de Médico Veterinario. Cutervo, Perú: Universidad Nacional Pedro Ruiz Gallo, 2017.
- Andrade-Yucailla, Verónica; Mazo, Lida; Vargas, J. C. & Lima-Orozco, R. Comportamiento productivo de cuyes en crecimiento-ceba alimentados con forraje de *Ipomoea batatas* L. en la región Amazónica Ecuatoriana. *UTCiencia Ciencia y Tecnología al servicio del pueblo*. 2 (1):24-28, 2015.
- AOAC. *Official methods of analysis*. 18th ed. Maryland, USA: Association of Official Agricultural Chemist, 2016.
- Avalos-Sánchez, Consuelo del R. *Utilización de la caña de azúcar fresca y picada (20, 40, 60 y 80 %) más alfalfa en crecimiento y engorde de cuyes*. Trabajo de grado presentado como requisito para optar el título profesional de Ingeniero Zootecnista. Riobamba, Ecuador: Escuela Superior Politécnica de Chimborazo, 2010.
- Bazán-García, A.; Cubas-Bazan, P.; Canches-Gonzales, A.; Díaz-García, América & Mariano-Santiago, H. Parámetros productivos en cuyes de la línea Perú (*Cavia porcellus*) alimentados con alfalfa fresca (*Medicago sativa*) y diferentes proporciones de harina de sangre de bovino. *Investigación Valdizana*. 10 (1):21-24, 2016.
- Camino, J. & Hidalgo, V. Evaluación de dos genotipos de cuyes (*Cavia porcellus*) alimentados con concentrado y exclusión de forraje verde. *Rev. investig. vet., Perú*. 25 (2):190-197, 2014.
- Canto, F.; Bernal, W. & Saucedo, J. Efecto de suplementación con probiótico (*lactobacillus*) en dietas de alfalfa y concentrado sobre parámetros productivos de cuyes mejorados en crecimiento y engorde. *Revista Científica UNTRM: Ciencias Naturales e Ingeniería*. 2 (2):39-44, 2018.
- Castro-Bedriñana, J.; Chirinos-Peinado, Doris & Calderón-Inga, J. Calidad nutricional del rastrojo de maca (*Lepidium peruvianum* Chacón) en cuyes. *Rev. investig. vet., Perú*. 29 (2):410-418, 2018. DOI: <http://dx.doi.org/10.15381/rivep.v29i2.13405>.
- Choez, Katherine & Ravillet, V. Frejol castilla (*Vigna unguiculata* L. Walp) como ingrediente en raciones de crecimiento-engorde de cuyes (*Cavia porcellus*) mejorados. *Rev. investig. vet., Perú*. 29 (1):180-187, 2018. DOI: <http://dx.doi.org/10.15381/rivep.v29i1.14086>.
- Gobierno Provincial del Azuay. *Plan de desarrollo y ordenamiento territorial del Azuay 2015-2030*. No. 12. Cuenca, Ecuador: Dirección de Planificación y Coordinación de Ordenamiento Territorial, 2015.
- Gualoto-Lata, G. A. *Evaluación de diferentes niveles de harina de Pennisetum violaceum (maralfalfa) en la elaboración de bloques nutricionales y su utilización en la alimentación de cuyes en la etapa de crecimiento y engorde*. Trabajo de grado presentado como requisito para optar por el título de Ingeniero Zootecnista. Riobamba, Ecuador: Escuela Superior Politécnica de Chimborazo, 2018.
- Guevara, J. & Carcelén, F. Efecto de la suplementación de probióticos sobre los parámetros productivos de cuyes. *Rev. Per. Quím. Ing. Quím.* 17 (2):69-74, 2014.
- Meza-Bone, G. A.; Cabrera-Verdezoto, R. P.; Morán-Morán, Jéssica J.; Meza-Bone, F. F.; Cabrera-Verdesoto, C. A.; Meza-Bone, C. J. et al. Mejora de engorde de cuyes (*Cavia porcellus* L.) a base de gramíneas y forrajeras arbustivas tropicales en la zona de Quevedo, Ecuador. *Idestia (Arica)*. 32 (3):75-80, 2014. DOI: <http://dx.doi.org/10.4067/S0718-34292014000300010>.
- Paillacho-Sánchez, W. R. *Evaluación de una dieta a base de harina de yuca (Manihot esculenta) y de alfalfa (Medicago sativa) en un balanceado para la*

- alimentación de cuyes (*Cavia aperea porcellus*, L.) en la etapa de engorde. Trabajo de titulación previa la obtención del título de Ingeniero en Desarrollo Integral Agropecuario. Tulcán, Ecuador: Universidad Politécnica Estatal del Carchi, 2017.
- Rivera, J. E.; Chará, J.; Gómez-Leyva, J. F.; Ruíz, T. & Barahona, R. Variabilidad fenotípica y composición fitoquímica de *Tithonia diversifolia* A. Gray para la producción animal sostenible. *LRRD*. 30 (2). <http://www.lrrd.org/lrrd30/12/rive30200.html>, 2018.
- Rosales-Jaramill, C. A.; Rodas-Carpi, E. R.; Nieto-Escandón, P. E.; Torres-Inga, C. S.; Gordillo-Guambana, B. G.; Aucapiña, C. *et al.* Extirpación de las espículas del pene de cuy (*Cavia porcellus*) y su efecto sobre la ganancia de peso y agresividad. *Rev. prod. anim.* 30 (1):47-52, 2018.
- Salazar-Ortiz, J.; Trejo-Téllez, L. I.; Valdez-Balero, A.; Senties-Herrera, H. E.; Rosas-Rodríguez, M.; Gallegos-Sánchez, J. *et al.* Caña de azúcar (*Saccharum* spp.) en la alimentación de rumiantes: experiencias generadas con cañas forrajeras. *Agroproductividad*. 10 (11):70-75, 2017.
- Sánchez-Laiño, A.; Sánchez-Gallardo, S.; Godoy-Becerra, S.; Díaz-Ocampo, R. & Vega-Pastuña, Norma. Gramíneas tropicales en el engorde de cuyes mejorados sexados (*Cavia porcellus* Linnaeus) en la zona de la maná. *Ciencia y Tecnología*. 2:25-28, 2009.
- Toro-Molina, Blanca; Yáñez-Zapata, M.; Andrade-Aulestia, Patricia; Labrada-Ching, J.; Chacón-Marcheco, E.; Zambrano-Cuadro, Natalia *et al.* La inclusión del bagazo de caña en la ración de cuyes (*Cavia porcellus*) de engorde. *REDVET. Rev. Electrón. vet.* 18 (10):1-6. <https://www.redalyc.org/articulo.oa?id=63653470030>, 2017.
- Valenzuela-Rocha, Rocío. *Determinación de la digestibilidad y energía digestible del forraje seco de mucuna (*Stizolobium deeringianum*) en cuyes*. Tesis para optar por el título de Ingeniero Zootecnista. Lima: Facultad de Zootecnia, Universidad Nacional Agraria La Molina, 2015.
- Vivas-Quila, N. J.; Criollo-Dorado, Milvia & Cedeno-Gómez, María C. Frecuencia de corte de pasto elefante morado *Pennisetum purpureum* Schumacher. *Rev. Bio. Agro.* 17 (1):45-55, 2019. DOI: <http://dx.doi.org/10.18684/bsaa.v17n1.1203>.
- Yamada, Graciela; Bazán, V. & Fuentes, Nadia. Comparación de parámetros productivos de dos líneas cárnicas de cuyes en la costa central del Perú. *Rev. investig. vet., Perú.* 30 (1):240-246, 2019. DOI: <http://dx.doi.org/10.15381/rivep.v30i1.15678>.